

Erratum: Glassy Chimeras Could Be Blind to Quantum Speedup... [Phys. Rev. X 4, 021008 (2014)]

Martin Weigel,¹ Helmut G. Katzgraber,^{2,3,4} Jonathan Machta,^{5,4} Firas Hamze,⁶ and Ruben S. Andrist⁴
(The Octomore Collaboration)

¹Applied Mathematics Research Centre, Coventry University, Coventry, CV1 5FB, England

²Department of Physics and Astronomy, Texas A&M University, College Station, Texas 77843-4242, USA

³Materials Science and Engineering Program, Texas A&M University, College Station, Texas 77843, USA

⁴Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

⁵Physics Department, University of Massachusetts, Amherst, MA 01003 USA

⁶D-Wave Systems, Inc., 3033 Beta Avenue, Burnaby, British Columbia, V5G 4M9, Canada

(Dated: December 22, 2014)

PACS numbers: 75.50.Lk, 75.40.Mg, 05.50.+q, 03.67.Lx




FIG. 1: Log-log plot of the FM susceptibility at its peaks, $\chi_m(\beta = \beta_{\max})$, as a function of the effective linear system size $\sqrt{N}$. The data fall onto a straight line, showing that corrections to scaling are very small. The dashed line has slope $2 - \eta = 7/4$.

In Phys. Rev. X 4, 021008 (2014), Katzgraber *et al.* studied Ising spins on the Chimera topology [1] of the D-Wave Two quantum annealing machine both in the spin-glass (SG), as well as the ferromagnetic (FM) sector. For the simulations in the FM sector, the following critical parameters based on simulations of systems of up to $N = 3200$ spins were computed: $\beta_c^m = 0.2402(3)$, $\nu_m \approx 1$, and $\eta_m \approx 2/5$. Here, $\beta_c^m = 1/T_c^m$ is the inverse critical temperature, ν_m the critical exponent of the FM correlation length, and η_m the critical exponent associated with the FM susceptibility. Using single-cluster updates [2], Monte Carlo simulations of up to $N = 131\,072$ spins in the FM sector have been performed. A detailed finite-size scaling analysis of the specific heat, the magnetization, the susceptibility, the Binder parameter and related cumulants, as well as three logarithmic deriva-

tives of the magnetization together with an analysis of cross-correlations [3, 4] leads to the following estimates of the critical parameters: $\beta_c^m = 0.2411929(26)$, $\nu_m = 0.9980(15)$, and $\eta_m = 0.2513(14)$. Hence, the critical exponents of a FM Ising model on the Chimera lattice are in agreement with the exact values for the two-dimensional (2D) Ising model, namely $\nu_{2D} = 1$ and $\eta_{2D} = 1/4$ [5]. Therefore, both models share the same universality class [6]. Figure 1 illustrates the perfect agreement between our new simulations and the exact values. Plotted are the FM susceptibility at its maximum, i.e., for $\beta = \beta_{\max}(N)$, vs the number of spins, where $\chi_m(\beta = \beta_{\max}) \sim (\sqrt{N})^{2-\eta_m}$. Indeed, the dashed line has slope $2 - \eta_{2D} = 7/4$ and is a guide to the eye. We would like to conclude by emphasizing that

1. For the FM sector, only the critical exponent η_m is incorrect in Ref. [7], as well as the claim that the universality class of FM Ising models in 2D and on the Chimera topology might be different.
2. For the SG sector $\eta_q = 0$ or close to zero is plausible. However, simulations to verify this would be difficult.
3. The main conclusions of Ref. [7] remain unchanged.
4. We would like to caution researchers seeking quantum speedup based on systems with 512 qubits or less, because, while corrections to scaling are weak, they seem to persist up to large system sizes.

Acknowledgments — M.W. and H.G.K. acknowledge support from the European Commission through the IRSES network DIONICOS under contract No. PIRSES-GA-2013-612707. H.G.K. acknowledges support from the NSF (Grant No. DMR-1151387) and would like to thank Bruichladdich Dist. for providing continued inspiration. J.M. acknowledges support from NSF (Grant No. DMR-1208046).

[1] P. Bunyk, IEEE Transactions on Applied Superconductivity **24**, 1700110 (2014).
[2] U. Wolff, Phys. Rev. Lett. **62**, 361 (1989).
[3] M. Weigel and W. Janke, Phys. Rev. Lett. **102**, 100601 (2009).
[4] M. Weigel and W. Janke, Phys. Rev. E **81**, 066701 (2010).
[5] J. M. Yeomans, *Statistical Mechanics of Phase Transitions* (Ox-

ford University Press, Oxford, 1992).
[6] Yes, Peter, you were right...
[7] H. G. Katzgraber, F. Hamze, and R. S. Andrist, Phys. Rev. X **4**, 021008 (2014).